

## Cumulative Index 2005

*Note:* Page numbers of article titles are in **boldface** type.

### A

Ablation, tissue, multilevel radiofrequency, for treatment of obstructive sleep apnea, 531

Acetaminophen, as adjuvants to intravenous patient-controlled analgesia, 113  
for postoperative pain in children, 165–166  
in multimodal approach to postoperative pain management, 188–189  
nitric oxide-releasing derivatives of, 191

Acupuncture, in the preoperative period, impact on child's anxiety, 605–606

Acute lung injury, transfusion-related, 247

Acute normovolemic hemodilution, **271–281**  
benefits of, 277  
cost effectiveness of, 278  
efficacy of, 277–278  
future of, 278  
in cardiac surgery, 340  
in elective orthopedic surgery, 308  
in pediatric patients, 350–353  
physiologic consequences, 273–274  
practical considerations for, 275–277  
technique, 271–273

Acute pain nurses, 216–221

Acute pain service, **211–225**  
audits and continuous quality improvement of, 221  
cost-effectiveness of, 215–216  
impact on outcome, 215  
implementation of, 216–221  
education, 219–220  
role of surgeon in, 219  
specialist pain nurse-based, 220–221  
upgrading role of ward nurses in, 217–218  
prevalence of, 212–213  
structure and function of, 213–214  
Adenotonsillectomy, in children, for obstructive sleep apnea treatment, 539–540

Adjuncts, analgesic, for postoperative pain management, **85–107**

alpha-2 agonists, 90–93  
dexmedetomidine, 91, 94  
neuraxial clonidine, 91  
peripheral nerve blocks with clonidine, 91  
systemic clonidine, 90–91  
anticonvulsants, 94–95  
corticosteroids, 98  
epidural administration of, 131  
in children, 178–180

alpha-2-agonists, 179  
epinephrine, 178  
ketamine, 179–180  
neostigmine, 180  
neuraxial opioids, 179  
sodium bicarbonate, 179  
tramadol, 180

N-methyl-D-aspartate (NMDA)  
antagonists, 85–90  
amantadine, 90  
dextromethorphan, 88–89  
ketamine, 86–88  
magnesium, 89  
neostigmine, 96–98  
opioid antagonists, 95–96

Adjustable gastric banding, laparoscopic, management of obesity with, 509–511

Administration, of blood, errors in, in transfusion medicine, 257–258

Adolescents, ethical issues in pediatric palliative care, 851  
obese, bariatric surgery in, 756–757

Advance care planning, in pediatric palliative care, 849–850

Age, as risk factor for obstructive sleep apnea, 412–413

Air embolism, complication of red cell recovery and reinfusion, 290

Airway anatomy, in children, pathophysiology of obstructive sleep apnea and, 536

- Airway assessment, preoperative, in obese patients, 465-466
- Airway complications, after pediatric lung transplantation, 725
- Airway injury, avoidance of, during single-lung ventilation for pediatric anesthesia, 704
- Airway, difficult, rapid sequence induction and, in patients with obesity and/or obstructive sleep apnea, 552-555
- Alcohol consumption, as risk factor for obstructive sleep apnea, 414-415
- Alfentanil, in pediatric anesthesia, 628
- Allogeneic blood transfusion, in cardiac surgery patients, clinical outcomes in, 334-339
- Alpha-2 adrenergic agonists, pediatric acute pain management with, 798
- Alpha-2 agonists, as adjunct for postoperative pain in children, 179  
as analgesic adjunct in postoperative pain management, 90-93  
in multimodal approach, 192-193
- Amantadine, as analgesic adjunct in postoperative pain management, 90
- Anemia, in surgical patients, clinical outcomes of, **315-325**  
efficacy of transfusion, 319-322  
human data, 318-319  
interaction between pathophysiologic processes and, 317-318  
pathophysiology, 315-317  
transfusion threshold, 322-323
- Anesthesia, in obese patients, **479-486**  
intensity of monitoring required for, 483  
laparoscopy in, 484  
minimizing hypoxia during anesthesia, 484-486  
pharmacokinetics, 479-481  
positioning obese patients for surgery, 481-482  
regional anesthesia in, 482-483  
in patients with obstructive sleep apnea, in children, 542-544  
effects of anesthetic drugs on ventilation responses in, 486-487  
intensity of intraoperative monitoring required, 487  
rational approach to, 488  
technique, 487  
pediatric, blood conservation in, **347-361**  
acute normovolemic hemodilution, 350-353  
deliberate hypotension, 354  
human recombinant erythropoietin, preoperative administration of, 354-356  
intraoperative blood salvage, 353  
multimodal strategies for, 356-357  
preoperative autologous donation, 349-350  
transfusion triggers and institutional algorithms, 348-  
pediatric. *See Pediatric anesthesia.*  
rapid sequence induction in patients with obesity and/or obstructive sleep apnea, **551-564**  
cricoid pressure, 556-560  
induction agent pharmacology, 560-561  
risk of gastroesophageal reflux and pulmonary aspiration, 555-556  
with the difficult airway, 552-555
- Anesthesia ventilators. *See Ventilatory support, mechanical.*
- Anesthetic agents, new developments in pediatric cardiac anesthesia, 655-658  
local. *See Local anesthetics.*
- Anticoagulant, in red cell recovery and reinfusion, 286-287
- Anticonvulsants, as analgesic adjuncts in postoperative pain management, 94-95
- Antiemetics, as adjuvants to intravenous patient-controlled analgesia, 113-114
- Antifibrinolytic agents, in pediatric cardiac anesthesia, 658-659, 659-660
- Antispasmodics, pediatric acute pain management with, 799-800
- Anxiety, preoperative, in children, psychological preparation for, **597-614**  
risk factors for development of, 598-599  
treatment of, 599-606  
acupuncture, 605-606  
behavioral intervention, 606  
music, 604-605  
outcomes of, 606-609  
parental presence during induction of anesthesia, 601-604  
preoperative preparation programs, 599-601
- Apnea-hypopnea index, 407, 446-447

Apnea. See also *Obstructive sleep apnea*.  
definition of, 406-407

Arndt endobronchial blockers, for single-lung ventilation in pediatric anesthesia, 698-700

Arrhythmias, after pediatric lung transplantation, 726

Artificial oxygen carriers, current status of, 373-389

- hemoglobin solutions, 375-382
  - diaspirin cross-linked hemoglobin, 376-378
  - hemoglobin raffiner, 380-381
  - human polymerized hemoglobin, 379-380
  - human recombinant hemoglobin, 378
  - maleimide-activated polyethylene glycol-modified hemoglobin, 381-382
  - polymerized bovine hemoglobin-based, 378-379
- nano-dimension artificial red blood cells, 384
- outlook for, 384-385
- perfluorocarbon (PFC) emulsions, 382-384
- perflubron emulsion (Oxygent), 382-384

Aspiration, pulmonary, risk of, in anesthesia of patients with obesity and/or obstructive sleep apnea, 555-556

Assessment, preoperative.

See *Preoperative evaluation*.

Atherosclerotic cardiovascular disease, associated with obstructive sleep apnea, 439

Autologous blood donation

- acute normovolemic hemodilution, 271-281, 308
- benefits of, 277
- cost effectiveness of, 278
- efficacy of, 277-278
- future of, 278
- physiologic consequences, 273-274
- practical considerations for, 275-277
- technique, 271-273
- for elective orthopedic surgery, perioperative assessment of patient, 295-303
- preoperative evaluation and methods to reduce blood use, 305-313

preoperative, 263-270

- aggressive phlebotomy in, 266-267
- cost effectiveness, 268
- erythropoietin use in, 266-267
- in pediatric patients, 349-350
- patient selection for, 264-266
- preoperative collection of components, 267-268
- transfusion trigger in, 267
- red blood cell recovery and reinfusion, 283-294, 308-309

Axillary block, in children, 822-824

Axillary brachial plexus block, for postoperative pain, 145-146

## B

B19 parvovirus infection, transfusion-transmitted, 245

Baby Doe, ethical issues in pediatric palliative care, 850

Bacterial contamination, transfusion-transmitted infection due to, 244

Balloon-tipped bronchial blockers, for single-lung ventilation in pediatric anesthesia, 697

Banding, gastric, laparoscopic adjustable, management of obesity with, 509-511

Barbiturates, for procedural sedation in pediatric patients, 645, 648

Bariatric surgery, 504-520

- choice of surgery, 518-519
- complications, 517-518
- future developments, 517
- in obese adolescents, 756-757
- indications for, 505
- malabsorptive procedures, 514-516
  - biliopancreatic diversion, 514-516
  - duodenal switch, 514-516
- overview of surgical procedures, 506-508
- preoperative preparation, 505
- restrictive procedures, 508-514
  - laparoscopic adjustable gastric banding, 509-511
  - roux-en-Y gastric bypass, 511-514
  - vertical gastric banding, 508-509
  - simultaneous cholecystectomy with, 519

Barriers, to health care, in pediatric palliative care, 851-853

Behavioral consequences, of inadequate postoperative pain relief, 23-24

Behavioral interventions, in the preoperative period, impact on child's anxiety, 605-606

- Behavioral modification, in obesity management, 502  
in obstructive sleep apnea management, 527
- Benzodiazepines, for procedural sedation in pediatric patients, 645
- Biliopancreatic diversion, surgical management of obesity with, 514-516
- Biology, of obesity and obstructive sleep apnea, **445-461**  
definitions, thresholds, and methodology, 446-447  
familial relationships in, 447-449  
mechanisms linking genetic variability in, 451-453  
obesity and, 449-451  
other inherited phenotypes predisposing to, 453-457  
abnormalities in sleep regulation, 456-457  
alterations in ventilation control, 455-456  
craniofacial morphology, 453-455  
specific genes and genetic markers associated with, 449
- Bleeding, new developments in pediatric cardiac anesthesia, 658-659  
perioperative, in cardiac surgery, 327-329
- Blood conservation, in cardiac surgery, **327-345**  
in critically ill patients, **363-372**  
in orthopedic surgery, **305-313**  
in pediatric anesthesia, **347-361**  
with artificial oxygen carriers, **373-389**
- Blood donation, autologous, **263-270**  
acute normovolemic hemodilution, **271-281**  
aggressive phlebotomy in, 266-267  
cost effectiveness, 268  
erythropoietin use in, 266-267  
patient selection for, 264-266  
preoperative collection of components, 267-268  
transfusion trigger in, 267
- Blood loss, during cell salvage, calculation of, 288
- Blood transfusions. *See also Transfusion medicine.*  
risks of, **241-252**  
acute lung injury related to, 247  
emerging risks, 248  
errors in, 247-248, **253-261**  
immunosuppression, 246-247  
infections, 241-246
- Bone healing, effects of cyclooxygenase-2 inhibitors for postoperative pain management, 61-62
- Bovine hemoglobin-based oxygen carrier, polymerized, 378-379
- Brachial plexus blocks, for postoperative pain, 141-146  
axillary, 145-146  
in children, 822-824  
infraclavicular, 144-145  
interscalene, 141-143  
supraclavicular, 143-144
- Bronchial blockers, balloon-tipped, for single-lung ventilation in pediatric anesthesia, 697
- Bronchiolitis obliterans, after pediatric lung transplantation, 724-725
- Bupivacaine, for postoperative pain in children, 176-177
- 
- ## C
- Cancer, risks of in obesity, 427
- Cardiac anesthesia, pediatric, new developments in, **655-676**  
anesthetic agents, 655-658  
bleeding and coagulation, 658-662  
mechanical circulatory support, 666-667  
neurological monitoring, 662-666  
short-term support, 667-672
- Cardiac surgery, conventional *versus* bloodless, **327-345**  
blood conservation methods in, 339-341  
clinical outcomes, factors influencing, 334-339  
coronary artery bypass graft techniques and, 330  
laboratory coagulation tests, 330-331  
predictors of transfusion requirements for, 332-334  
variability in transfusion practice for, 331-332
- Cardiorenal effects, of cyclooxygenase-2 inhibitors for postoperative pain management, 61-62
- Cardiovascular disease, associated with obstructive sleep apnea, 439  
risk in patients with obstructive sleep apnea, preoperative evaluation of, 473
- Cardiovascular pathophysiology, in obesity, 422

- Cardiovascular system, preoperative assessment in obese patients, 467-469
- Catheter techniques, for delivery of local anesthetics for postoperative pain management, 76-77
- Catheters, Fogarty embolectomy, for single-lung ventilation in pediatric anesthesia, 697-698
- Caudal analgesia, for pediatric acute pain management, 807  
in children, 170-175  
    continuous, 173-174  
    high blocks, 172-173  
    single-shot, 171-172
- Caudal anesthesia, in children, 824-827
- Celecoxib, efficacy in postoperative pain management, 53-54
- Cell salvage, red blood cell recovery and reinfusion, **283-294**  
    comparison with other techniques, 290-292  
    cell salvage, 292  
    normovolemic hemodilution, 291-292  
    preoperative autologous donation, 291  
    complications, 290  
        air embolism, 290  
        wrong wash solution, 290  
    in elective orthopedic surgery, 308-309  
    in pediatric patients, 353  
    optimizing red cell return, anticoagulant, 287  
        calculation of blood loss during cell salvage, 288  
    collection reservoir, 287-288  
    optimizing red cell return, 285-288  
    rinsing of surgical sponges, 286  
    suction, 285-286  
    wash quality, 289-290  
    physics of cell salvage machine, 283-285
- Central sleep apnea, 407-408, 431
- Centroneuraxis opioids, for pediatric acute pain management, 808
- Cerebral palsy, surgery and anesthesia for children who have, **733-743**  
    anesthetic management, 735-742  
        bleeding, intraoperative, 740-742  
        intraoperative, 735-736, 739  
        preoperative, 735  
        regional techniques, 739  
        temperature, 739-740  
    preoperative concerns, 734-735  
        gastrointestinal, 734  
        neurological, 734-735  
        respiratory, 734
- Cerebrovascular disease, associated with obstructive sleep apnea, 439
- Chest. *See* Thoracic surgery.
- Children, anesthesia in.  
    *See* Pediatric anesthesia.  
    obesity in, 398-399  
    obstructive sleep apnea in, **535-549**  
        clinical features, 537-538  
        complications, 538  
        evaluation, 539  
        pathophysiology, 535-537  
            airway anatomy, 536  
            neuromotor factors, 537  
            obesity and, 536-537  
        treatment, 539-547  
            intraoperative anesthesia plan, 542-544  
            monitoring, 544-545  
            pain management, 546  
            perioperative management, 540  
            postoperative care, 544  
            preanesthesia evaluation, 540-542  
            prognosis, 546-547  
            respiratory support, 545-546  
            surgery for, 540  
    postoperative pain management in, **163-184**  
        epidural/caudal blocks, 170-175  
        intravenous patient/parent/nurse-controlled analgesia, 168-170  
        local anesthetics/adjuncts, 176-180  
        nonopioid analgesics, 165-168  
        potent opioid analgesics, 168
- Chloral hydrate, for procedural sedation in pediatric patients, 645
- Cholecystectomy.
- Cholecystectomy, laparoscopic, procedure-specific postoperative pain management guidelines for, 207-209  
    performed simultaneously with bariatric surgery, 519
- Chronic conditions, complex, palliative care in children with life-threatening, **837-856**
- Chronic opioid-consuming patients.  
    *See* Opioids.
- Chronic pain, postoperative. *See* Pain, postoperative, persistent.

- Clonidine, as analgesic adjunct in postoperative pain management, 90-93  
    neuraxial, 91  
    peripheral nerve blocks with, 91  
    systemic, 90-91
- Coagulation, new developments in pediatric cardiac anesthesia, 658-659
- Codeine, pediatric acute pain management with, 795-796
- Collection reservoir, in red cell recovery and reinfusion, 286-287
- Collection, of blood, errors in, in transfusion medicine, 257-258
- Communication, as part of pediatric palliative care, 843
- Complex chronic conditions, palliative care in children with, **837-856**
- Congenital heart disease, pediatric cardiac anesthesia, new developments in, **655-676**  
    anesthetic agents, 655-658  
    bleeding and coagulation, 658-662  
    mechanical circulatory support, 666-667  
    neurological monitoring, 662-666  
    short-term support, 667-672
- Congestive heart failure, associated with obstructive sleep apnea, 440
- Conservation, of blood, in cardiac surgery, **327-345**  
    in critically ill patients, **363-372**  
    in orthopedic surgery, **305-313**  
    in pediatric anesthesia, **347-361**  
    with artificial oxygen carriers, **373-389**
- Contamination, bacterial, transfusion-transmitted infection due to, 244
- Continuous epidural analgesia, pediatric acute pain management with, 803-805
- Continuous intravenous opioid infusion, pediatric acute pain management with, 803
- Continuous positive airway pressure (CPAP), for treatment of obstructive sleep apnea, 525-526
- Convalescence, postoperative, multimodal analgesia to shorten, 196
- Coronary artery bypass graft surgery.  
    See *Cardiac surgery*
- Coronary artery disease, associated with obstructive sleep apnea, 439
- Corticosteroids, as analgesic adjuncts in postoperative pain management, 98
- Cost effectiveness, of autologous blood donation, 268  
    of acute normovolemic hemodilution, 278
- Craniofacial abnormalities, inherited, predisposing to sleep apnea, 453-455
- Creutzfeld-Jakob disease, transfusion-transmitted, 245-246
- Cricoid pressure, rapid sequence induction and, in patients with obesity and/or obstructive sleep apnea, 555-560
- Critically ill patients, blood conservation in, **363-372**  
    red blood cell transfusions in, 363-66  
    efficacy and risks, 364-366
- Cyclooxygenase-2 (COX-2) inhibitors, in postoperative pain management, contraindications, 63-64  
    drug interactions, 63  
    efficacy, 51-57  
        celecoxib, 53-54  
        parecoxib, 56-57  
        valdecoxib, 55-56  
    expression and function, 50-51  
    in children, 167  
    in multimodal approach, 189-191  
    safety and tolerability, 57-63  
        cardiorenal effects, 57-61  
        effects on bone and wound healing, 62-63  
        gastrointestinal toxicity, 57  
        hematologic effects, 57-58  
        hepatic effects, 61-62
- Cytomegalovirus infection, transfusion-transmitted, 244-245
- 
- D**
- Death, developmental stages of children's understanding of, 839-842  
    palliative care in pediatrics, **837-856**
- Delivery techniques, novel, for local anesthetics in postoperative pain management, 75-77  
    catheter techniques, 76-77  
    encapsulated drugs, 75-76
- Demographics, of obesity, **397-403**  
    of obstructive sleep apnea, **405-420**
- Development, child, impact of complex chronic conditions on, 839-842
- Developmental pharmacology, and opioids in pediatric anesthesia, 622-625

- Dexmedetomidine, as analgesic adjunct in postoperative pain management, 91, 94  
pediatric acute pain management with, 798
- Dextromethorphan, as analgesic adjunct in postoperative pain management, 88–89  
for postoperative pain in children, 167
- Diabetes, associated with obstructive sleep apnea, 438–439
- Diagnosis, of obesity, **397–403**  
of obstructive sleep apnea, **405–420**  
during preoperative evaluation, 470–472
- Diaspirin cross-linked hemoglobin, as artificial oxygen carrier, 376–378
- Diet, in obesity management, 502
- Disordered breathing events, in obstructive sleep apnea, 406–408
- Donation, blood. *See* Blood donation.
- Double-lumen tubes, for single-lung ventilation in pediatric anesthesia, 701–703
- Drug clearance, developmental pharmacology and opioids in pediatric anesthesia, 622–625
- Drugs, for weight loss, in management of obesity, 502–504
- Duodenal switch, surgical management of obesity with, 514–516
- Dying, palliative care in pediatrics, **837–856**

## E

- Echocardiography, in pediatric cardiac anesthesia, measurement of response, 657–658
- Education, patient, in acute pain service, 219–220  
on intravenous patient-controlled analgesia, 116–117
- Elderly patients, intravenous patient-controlled analgesia in, 118–119
- Elective surgery, orthopedic, perioperative assessment of patients, **295–303**  
preoperative evaluation and methods to reduce blood use in, **305–313**
- Electroencephalogram modalities, in pediatric cardiac anesthesia, 662–663
- Embolectomy catheters, Fogarty, for single-lung ventilation in pediatric anesthesia, 697–698
- Embolism, air, complication of red cell recovery and reinfusion, 290
- Encapsulated drugs, for delivery of local anesthetics in postoperative pain management, 75–76
- Endobronchial blockers, Arndt, for single-lung ventilation in pediatric anesthesia, 698–700
- Endotracheal tubes, for single-lung ventilation in pediatric anesthesia, 697
- Epidemiology, of life-threatening illness in children, palliative care and, 837–839  
of obstructive sleep apnea, 494
- Epidural analgesia, for postoperative pain, **125–140**  
benefits and patient outcomes, 126–128  
continuous infusion *versus* patient-controlled, 132  
duration of, 131–132  
factors affecting efficacy of, 129–132  
in children, 170–175  
continuous, 173–175  
patient-controlled, 175–176  
in chronic opioid-consuming patients, 39  
risks of, 132–134  
in multimodal approach to postoperative pain management, 187  
pediatric acute pain management with, continuous, 803  
patient-controlled, 805–806
- Epidural hematoma, risk of, with epidural analgesia, 132–133
- Epidural opioids, in pediatric anesthesia, 630–632, 828–830
- Epinephrine, as adjunct for postoperative pain in children, 178
- Epoetin alfa, reducing blood use in elective orthopedic surgery with, 309–311
- Errors, medical, in transfusion medicine, 247–248, **253–261**  
prevention of, 258–259  
types of, 255–258  
UK Serious Hazards of Transfusion (SHOT) scheme, 254–255
- Erythropoietin, blood conservation with, in autologous blood donation, 266–267  
in cardiac surgery, 339–340  
in elective orthopedic surgery, 309–311  
in pediatric surgical patients, 354–356

- Ethical issues, in pediatric palliative care, 849-851
  - advance care planning, 849-850
  - Baby Doe, 850
  - in adolescents, 851

Ethnic differences, in obesity, 398, 401

Exercise, in obesity management, 502

## F

- Familial relationships, in sleep apnea, 447-449
- Family history, as risk factor for obstructive sleep apnea, 414
- Fascia iliacus compartment block, in children, 821
- Femoral nerve block, for postoperative pain, 148-150
  - in children, 819-821
- Fenfluramine, 503
- Fentanyl, in pediatric anesthesia, 626-627
  - pediatric acute pain management with, 794-795
- Fentanyl challenge, preoperative, in chronic opioid-consuming patients, 41
- Fogarty embolectomy catheters, for single-lung ventilation in pediatric anesthesia, 697-698

## G

- Gabapentin, as analgesic adjunct in postoperative pain management, 92-93, 94-95
  - in multimodal approach, 193
- Gastric banding, laparoscopic adjustable, management of obesity with, 509-511
- Gastric bypass, Roux-en-Y, surgical
  - management of obesity with, 511-514
  - complications following, 513
  - essential components of, 512
- Gastroesophageal reflux, risk of, in anesthesia of patients with obesity with, 511-514
  - complications following, 513
  - essential components of, 512
- Gastroesophageal reflux, risk of, in anesthesia of patients with obesity with, 511-514
  - complications following, 513
  - essential components of, 512
- Gastroesophageal reflux disease, associated with obstructive sleep apnea, 440
  - preoperative evaluation of obese patients with, 469
- Gastrointestinal complications, after pediatric lung transplantation, 725-726

Gastrointestinal toxicity, of cyclooxygenase-2 inhibitors for postoperative pain management, 57

Gender, as risk factor for obstructive sleep apnea, 412

Genetics, of malignant hyperthermia, 615-619
 

- of obesity and obstructive sleep apnea,

### 445-461

- definitions, thresholds, and methodology, 446-447
- familial relationships in, 447-449
- mechanisms linking genetic variability in, 451-453
- obesity and, 449-451
- other inherited phenotypes
  - predisposing to, 453-457
  - abnormalities in sleep regulation, 456-457
  - alterations in ventilation control, 455-456
  - craniofacial morphology, 453-455
  - specific genes and genetic markers associated with, 449
- role in obesity, 399-400

Genioglossus advancement, for treatment of obstructive sleep apnea, 530-531

Glucocorticoids, in multimodal approach to postoperative pain management, 193

Grief, as part of pediatric palliative care, 843

Guidelines, procedure-specific, for postoperative pain management, 203-210
 

- example of, for laparoscopic cholecystectomy, 207-209

## H

- Heart disease, congenital, pediatric cardiac anesthesia, new developments in, 655-676
  - anesthetic agents, 655-658
  - bleeding and coagulation, 658-662
  - mechanical circulatory support, 666-667
  - neurological monitoring, 662-666
  - short-term support, 667-672
- Heart transplantation, pediatric, 709-715
  - donor selection, 710-711
  - immunosuppressive agents, 713
  - indications, 709-710
  - intraoperative management, 711-712
  - outcome after, 713-715
  - recipient evaluation, 710
  - rejection, 715



- Hematologic effects, of cyclooxygenase-2 inhibitors for postoperative pain management, 57-58
- Hematoma, epidural, risk of, with epidural analgesia, 132-133
- Hemodilution. See *Acute normovolemic hemodilution*.
- Hemoglobin raffimer (Hemolink), 380-381
- Hemoglobin-based artificial oxygen carriers, 375-382  
  diaspirin cross-linked, 376-378  
  hemoglobin raffimer, 380-381  
  human polymerized hemoglobin, 379-380  
  human recombinant hemoglobin, 378  
  maleimide-activated polyethylene glycol-modified hemoglobin, 381-382  
  polymerized bovine hemoglobin-based, 378-379
- Hepatic drug clearance, developmental pharmacology and opioids in pediatric anesthesia, 622-625
- Hepatic effects, of cyclooxygenase-2 inhibitors for postoperative pain management, 61-62
- Hepatitis, transfusion-associated, 242-243
- History, of pediatric anesthesia, **573-595**  
  after World War II, 582-585  
  in the 1960s and 1970s, 585-587  
  in the 1980s and 1990s, 587-591  
  in the first half of the twentieth century, 578-582  
  in the nineteenth century, 575-578
- History, family, as risk factor for obstructive sleep apnea, 414
- Human immunodeficiency virus (HIV), transfusion-associated, 242-243
- Human recombinant hemoglobin, as artificial oxygen carrier, 378
- Human T-lymphotropic virus (HTLV)-I, transfusion-transmitted, 245
- Hydrocodone, pediatric acute pain management with, 796
- Hydromorphone, pediatric acute pain management with, 794
- Hyperalgesia, sensitization and, 2
- Hypertension, pulmonary, in patients with obstructive sleep apnea, preoperative evaluation of, 473  
  systemic, associated with obstructive sleep apnea, 438
- Hyperthermia, malignant, genetics of, **615-619**
- Hypopnea. See also *Obstructive sleep apnea*.  
  definition of, 406-407  
  physiologic characteristics of, 431-433
- Hypotension, deliberate, for blood conservation in pediatric surgical patients, 354
- Hypoxemia, treating and avoiding during single-lung ventilation for pediatric anesthesia, 703-704
- Hypoxia, during anesthesia, minimizing in obese patients, 484-486
- 
- I**
- Ilioinguinal-iliohypogastric nerve block, in children, 816-818
- Immunosuppression, due to blood transfusions, 246-247
- Immunosuppressive agents, for pediatric heart transplantation, 713
- Incisional pain, mechanisms of, **1-20**  
  clinical models of, 2-4  
  experimental *versus* clinical models of, 4  
  laparotomy models for, 4-5  
  plantar incision model for, 5-15  
  prevention strategies, 15-16  
  schematic for, 16-17
- Induction agents, pharmacology of, and rapid sequence induction in patients with obesity and/or obstructive sleep apnea, 560-561
- Infants, sensitivity to opioids in neonates, 624-625  
  spinal anesthesia for, 830-832  
    anatomy, 830-831  
    drug dosage in, 832  
    technique in ex-premature patients, 831
- Infection, after pediatric lung transplantation, 726  
  due to blood transfusions, 241-246  
    bacterial contamination, 244  
    cytomegalovirus, 244-245  
    other agents, 245-246  
      B19 parvovirus, 245  
      Creutzfeld-Jakob disease, 245-246  
      human T-lymphotropic virus, 245  
  risk of, with epidural analgesia, 133  
  viral, 242-243
- Infraclavicular brachial plexus block, for postoperative pain, 144-145

Interscalene brachial plexus block, for postoperative pain, 141-143

Intraoperative red blood cell retrieval and reinfusion. *See Cell salvage.*

Intravenous patient-controlled analgesia. *See Patient-controlled analgesia.*

## J

Jaw advancement techniques, for treatment of obstructive sleep apnea, 530

## K

Ketamine, as adjunct for postoperative pain in children, 179-180  
 as adjuvant to intravenous patient-controlled analgesia, 113  
 as analgesic adjunct in postoperative pain management, 86-88  
   role as adjunct to opioids, 86-88  
     as continuous bolus infusion, 87  
     as single intravenous bolus, 87-88  
   epidural, 88  
   in patient-controlled analgesia, 86-87  
   role in preventive analgesia, 86  
 pediatric acute pain management with, 797

## L

Laboratory errors, in transfusion medicine, 257

Laparoscopic adjustable gastric banding, surgical management of obesity with, 509-511

Laparoscopic cholecystectomy, procedure-specific postoperative pain management guidelines for, 207-209

Laparoscopy, in obese patients, 484

Laparotomy models, for postoperative pain, 4-5

Lateral decubitus position, physiology of, 693-695

Legal issues, in pediatric palliative care, 849-851

  advance care planning, 849-850  
   Baby Doe, 850  
   in adolescents, 851

Lepton, genetic variability in obesity and sleep apnea linked to, 451-453

Levobupivacaine, for postoperative pain in children, 177-178

Lingual muscles, physiotherapy of, for treatment of obstructive sleep apnea, 531

Liposomal drug delivery, for postoperative pain management, 75-76

Liposomes, hemoglobin-containing, 384

Local anesthetics, in postoperative pain management, 73-84

  beneficial actions, 79-81

  epidural, 130

    combined with opioids, 130-131

  in children, 176-178

    bupivacaine, 176-177

    newer agents, 177-178

  long-lasting, 77-78

  novel delivery techniques, 75-77

    catheter techniques, 76-77

    encapsulated drugs, 75-76

  shift to outpatient surgery and, 74-75

  toxicity issues, 78-79

  unexpected toxicity, 73-74

  pediatric acute pain management with, 798-799

  volume and dosage in children, 827-828

Local infiltration, in children, 824

Lumbar anesthesia, in children, 824-827

Lumbar epidural analgesia, continuous, in children, 174

Lumbar plexus block, for postoperative pain, 146-148

Lung injury, acute, transfusion-related, 247

Lung transplantation, pediatric, 715-728

  anesthesia management, 720-722

  contraindications for, 718-719

  history of, 715-716

  immunosuppression and surveillance, 723

  indications for, 717-718

  number of procedures, 716

  post-transplant considerations and complications, 723-727

  postoperative course and pain management, 722-723

  procedure, 719-720

  survival, 727-728

  waiting list, 716-717

## M

Magnesium, as analgesic adjunct in postoperative pain management, 89

- Malabsorptive procedures, for surgical management of obesity, 514-516  
biliopancreatic diversion, 514-516  
duodenal switch, 514-516
- Maleimide-activated polyethylene glycol-modified hemoglobin, 381-382
- Malignancy, after pediatric lung transplantation, 726-727
- Malignant hyperthermia, genetics of, **615-619**
- Mandibular repositioning devices, for treatment of obstructive sleep apnea, 526-527
- Markers, genetic, for obstructive sleep apnea, 449
- Mechanical ventilatory support, anesthesia ventilators, better options for children, 677-691  
in pediatric cardiac anesthesia, advances in, 666-667
- Meperidine, pediatric acute pain management with, 795
- Meridia. See *Sibutramine*.
- Metabolic syndrome, obesity and, 423-425
- Methadone, pediatric acute pain management with, 794
- Minimally invasive techniques, for treatment of obstructive sleep apnea, 530-531  
genioglossus advancement, 530  
multilevel radiofrequency tissue ablation, 531  
tongue-base suspension, 531
- Models, for postoperative pain, 2-15  
clinical, 2-4  
experimental *versus* clinical, 4  
laparotomy, 4-5  
plantar incision, 5-20
- Monitoring, during anesthesia, intensity of, for obese patients, 483  
for patients with obstructive sleep apnea, children, 544-545  
neurological, in pediatric cardiac anesthesia, 662-666
- Morbidly obese patients, intravenous patient-controlled analgesia in, 118
- Morphine, in pediatric anesthesia, 626  
pediatric acute pain management with, 792-794
- Multimodal analgesia, for postoperative pain management, postoperative and rehabilitation, **185-202**  
analgesic options for, 186-193  
as part of preemptive analgesia techniques, 195-196

optimal techniques for, 193-195  
to shorten convalescence, 196

Muscle strengthening, for treatment of obstructive sleep apnea, 531

Music, in the preoperative period, impact on child's anxiety, 604-605

Myocardial ischemia, risk in patients with obstructive sleep apnea, preoperative evaluation of, 473-474

## N

- N-methyl-D-aspartate (NMDA) antagonists, as adjunct in postoperative pain management, 85-90  
amantadine, 90  
dextromethorphan, 88-89  
in multimodal approach, 192  
ketamine, 86-88  
magnesium, 89
- Nalbuphine, pediatric acute pain management with, 796
- Naloxone, pediatric acute pain management with, 797
- Nano-dimension artificial red blood cells, 384
- Nasal surgery, for treatment of obstructive sleep apnea, 529
- Near-infrared spectroscopy, in pediatric cardiac anesthesia, 664
- Neonates. See also *Infants*.  
sensitivity to opioids, 624-625
- Neostigmine, as adjunct for postoperative pain in children, 180  
as analgesic adjunct in postoperative pain management, 96-98  
added to intravenous regional anesthesia, 98  
epidural, 97  
intra-articular, 97  
intrathecal, 96-97  
with local anesthetic for brachial plexus block, 97-98
- Nerve blocks, paravertebral. See *Paravertebral nerve blocks*.  
peripheral. See *Peripheral nerve blocks*.
- Neuraxial opioids, as adjunct for postoperative pain in children, 179
- Neurological monitoring, in pediatric cardiac anesthesia, 662-666
- Neuromotor factors, obstructive sleep apnea in children related to, 537

Nitric oxide, in derivatives of acetaminophen and NSAIDs, 191

Nonsteroidal anti-inflammatory drugs (NSAIDs), as adjuvants to intravenous patient-controlled analgesia, 112-113  
for postoperative pain in children, 166-167  
in multimodal approach to postoperative pain management, 188-189  
nitric oxide-releasing derivatives of, 191  
pediatric acute pain management with, 791-792

Normovolemic hemodilution. See *Acute normovolemic hemodilution*.

Nurses, role in acute pain service, specialist pain nurses, 220-221  
upgrading role of ward nurses in, 217-218

## O

Obesity, anesthetic management of patients with, **479-486**  
intensity of monitoring required for, 483  
laparoscopy in, 484  
minimizing hypoxia during anesthesia, 484-486  
pharmacokinetics, 479-481  
positioning obese patients for surgery, 481-482  
regional anesthesia in, 482-483  
biology and genetics of, **445-461**  
definitions and demographics, **397-403**  
ethnic and racial differences, 398, 401  
health costs associated with, 401-402  
in childhood, 398-399  
role of genetics in, 399-400  
socioeconomic factors, 401  
worldwide prevalence of, 397-398  
in children, obstructive sleep apnea and, 536-537  
morbid, intravenous patient-controlled analgesia in patients with, 118  
pathophysiologic changes of, **421-429**  
cancer, 427  
cardiovascular, 422  
metabolic syndrome, 423-425  
respiratory, 425-427  
pediatric, anesthesia for, **745-764**  
intraoperative, 758-759  
perioperative outcomes, 760-761

positioning, 759-760  
preoperative, 757-758  
bariatric surgery in adolescents, 756-757  
causes, 750-751  
definitions, 746  
pathophysiology related to, 751-754  
cardiovascular, 753-754  
endocrinology, 754  
respiratory, 752-753  
pharmacology related to, 754-756  
prevalence, 746-750  
postoperative considerations for patients with sleep apnea and, **493-500**  
preoperative evaluation of patients with, **463-478**  
airway assessment, 465-466  
cardiovascular system assessment, 468-469  
other considerations, 469-470  
respiratory function assessment, 466-467  
rapid sequence induction in patients with, **551-564**  
cricoid pressure, 556-560  
induction agent pharmacology, 560-561  
risk of gastroesophageal reflux and pulmonary aspiration, 555-556  
with the difficult airway, 552-555  
risk factor for obstructive sleep apnea, 411-412  
sleep apnea due to. See *Obstructive sleep apnea*.  
surgical and nonsurgical management of, **501-523**  
bariatric surgery, 504-520  
choice of surgery, 518-519  
complications, 517-518  
future developments, 517  
malabsorptive procedures, 514-516  
overview of surgical procedures, 506-508  
preoperative preparation, 505  
restrictive procedures, 508-514  
nonsurgical, 502-504  
behavioral modification, 502  
diet, 502  
exercise, 502  
weight loss drugs, 502-504

Obstructive sleep apnea, 397-564  
anesthetic management of patients with, **486-491**  
effects of anesthetic drugs on ventilation responses in, 486-487

- intensity of intraoperative monitoring required, 487
- rational approach to, 488
- technique, 487
- biology and genetics of obesity and, **445-461**
  - definitions, thresholds, and methodology, 446-447
  - familial relationships in, 447-449
  - mechanisms linking genetic variability in, 451-453
  - obesity and, 449-451
  - other inherited phenotypes predisposing to, 453-457
    - abnormalities in sleep regulation, 456-457
    - alterations in ventilation control, 455-456
    - craniofacial morphology, 453-455
  - specific genes and genetic markers associated with, 449
- demographics and diagnosis of, **405-420**
  - disease definition and diagnosis, 406-408
  - population prevalence, 408-410
  - risk factors for, 410-415
    - age, 412-413
    - alcohol consumption, 414-415
    - craniofacial abnormalities, 414
    - family history, 414
    - gender, 412
    - obesity, 411-412
    - race, 413
  - terminology, 406-407
- in children, **535-549**
  - clinical features, 537-538
  - complications, 538
  - evaluation, 539
  - pathophysiology, 535-537
    - airway anatomy, 536
    - neuromotor factors, 537
    - obesity and, 536-537
  - treatment, 539-547
    - intraoperative anesthesia plan, 542-544
    - monitoring, 544-545
    - pain management, 546
    - perioperative management, 540
    - postoperative care, 544
    - preanesthesia evaluation, 540-542
    - prognosis, 546-547
    - respiratory support, 545-546
    - surgery for, 540
- intravenous patient-controlled analgesia in patients with, 117-118
- obesity and. *See also Obesity:*
  - biology and genetics of, **445-461**
  - definitions and demographics, **397-403**
  - pathophysiologic changes of, **421-429**
- pathophysiology of, **431-443**
  - categories of, 431-433
  - causes, 434-435
  - contributing factors, 435-436
  - diseases associated with, 438-440
    - congestive heart failure, 440
    - coronary artery disease, 439
    - diabetes, 438-439
    - gastroesophageal reflux disease, 440
    - hypertension, 438
    - obesity, 438
    - stroke, 439
  - symptoms, 436-437
- postoperative considerations for patients with, **493-500**
  - additional considerations for obese patients, 498
  - disposition from postanesthesia care unit, 496-497
  - during recovery and immediate postoperative period, 495-496
  - pain management, 497-498
- preoperative evaluation of patients with, **463-478**
  - assessment of patients with known, 472-473
  - cardiovascular consequences of, 473-474
  - identifying patients with, 470-472
- rapid sequence induction in patients with, **551-564**
  - cricoid pressure, 556-560
  - induction agent pharmacology, 560-561
  - risk of gastroesophageal reflux and pulmonary aspiration, 555-556
  - with the difficult airway, 552-555
- treatment for, **525-534**
  - minimally invasive techniques, 530-531
    - genioglossus
      - advancement, 530
    - multilevel radiofrequency tissue ablation, 531
    - tongue-base suspension, 531
  - nonsurgical, 525-527
    - adjustable oral appliances, 526-527
    - continuous positive airway pressure, 525-526
    - weight loss and behavioral therapies, 527

- surgical, 527-532
    - jaw advancement techniques, 530
    - nasal surgery, 529
    - perioperative issues for, 531-532
    - physiotherapy of lingual and supra-hyoid muscles, 531
    - tonsillectomy, 528-529
    - tracheostomy, 529-530
    - uvulopharyngopalatoplasty, 527-528
  - Opioid antagonists, as analgesic adjuncts in postoperative pain management, 95-96
  - Opioids, efficacy of epidural analgesia with, 129-130
    - for postoperative pain in children, 168-176
      - epidural/caudal blocks, 170-176
      - patient-controlled, 175-176
    - intravenous patient/parent/nurse-controlled analgesia with, 168-170
    - in acute pediatric pain management, 792-797
      - centroneuraxis opioids, 808
      - continuous intravenous infusion, 803
    - in multimodal approach to postoperative pain management, 186-187
    - in pediatric anesthesia, 621-634
      - developmental pharmacology, 622-625
      - drugs commonly used in, 626-632
      - epidural, 828-830
      - for procedural sedation, 648
    - in postoperative care of chronic opioid-consuming patients, 37-48
      - defining threshold for respiratory depression, 41-42
      - intravenous patient-controlled analgesia with an opioid, 40-41, 117
      - practical considerations for, 44-45
      - predicting settings for, 42-43
      - transition to oral opioids, 46
    - preoperative fentanyl challenge, 41
    - preoperative plan for, 38-39
    - regional analgesia techniques and nonopioid analgesics, 39-40
    - epidural analgesia and peripheral nerve blocks, 39
    - nonopioid analgesics, 39-40
    - tolerance, 38
    - neuraxial, as adjunct for postoperative pain in children, 179
  - Oral analgesics, transition to, for pediatric acute pain management, 806-807
  - Oral appliances, for treatment of obstructive sleep apnea, 526-527
  - Organ transplant surgery, pediatric solid organ transplant, anesthesia considerations for, 709-731
    - heart transplantation, 709-715
    - lung transplantation, 715-728
  - Orlistat, in management of obesity, 503-504
  - Orthopedic elective surgery, perioperative assessment of patients, 295-303
    - preoperative evaluation and methods to reduce blood use in, 305-313
  - Outcomes, clinical, of surgical patients requiring transfusion, anemia and, 315-325
    - in cardiac surgery, 327-345
  - Outpatient surgery, shift to, and role of local anesthetics, 74-75
  - Oxycodone, pediatric acute pain management with, 796
  - Oxygen carriers, artificial. See *Artificial oxygen carriers*.
- 
- P**
- Pain management, in pediatric palliative care, 844-849
    - in single-lung ventilation for pediatric anesthesia, 705
  - opioids in pediatric anesthesia, 621-634
    - pediatric acute, 789-814
      - caudal analgesia, 807
      - centroneuraxis opioid, 808
      - continuous epidural analgesia, 803-805
      - cutaneous intravenous opioid infusion, 803
      - developmental neurobiology, 790
      - developmental pharmacology, 790-791
      - developmentally appropriate pain assessment, 800-801
      - drugs for, 791-800
      - nonpharmacologic methods, 809
      - patient-controlled analgesia, 801-803
        - epidural, 805-806
      - peripheral nerve blocks, 808-809
      - postoperative strategies, 801
      - regional techniques, 807
      - transition to oral analgesics, 806-807

- postoperative, 1-225
- acute pain service for, **211-225**
    - audits and continuous quality improvement of, 221
    - cost-effectiveness of, 215-216
    - impact on outcome, 215
    - implementation of, 216-221
    - prevalence of, 212-213
    - structure and function of, 213-214
  - analgesic adjuncts in, **85-107**
    - alpha-2 agonists, 90-93
      - dexmedetomidine, 91, 94
      - neuraxial clonidine, 91
      - peripheral nerve blocks with clonidine, 91
      - systemic clonidine, 90-91
    - anticonvulsants, 94-95
    - corticosteroids, 98
    - N-methyl-D-aspartate (NMDA) antagonists, 85-90
      - amantadine, 90
      - dextromethorphan, 88-89
      - ketamine, 86-88
      - magnesium, 89
    - neostigmine, 96-98
    - opioid antagonists, 95-96
  - cyclooxygenase-2 inhibitors in, **49-72**
    - contraindications, 63-64
    - drug interactions, 63
    - efficacy, 51-57
      - celecoxib, 53-54
      - parecoxib, 56-57
      - valdecoxib, 55-56
    - expression and function, 50-51
    - safety and tolerability, 57-63
      - cardiorenal effects, 57-61
      - effects on bone and wound healing, 62-63
      - gastrointestinal toxicity, 57
      - hematologic effects, 57-58
      - hepatic effects, 61-62
  - epidural analgesia for, **125-140**
    - benefits and patient outcomes, 126-128
    - factors affecting efficacy of, 129-132
    - risks of, 132-134
    - for patients with obstructive sleep apnea, 497-498
    - children, 546
  - in children, **163-184**
    - epidural/caudal blocks, 170-175
    - intravenous patient/parent/nurse-controlled analgesia, 168-170
    - local anesthetics/adjuncts, 176-180
    - nonopioid analgesics, 165-168
    - potent opioid analgesics, 168
  - in chronic opioid-consuming patients, **37-48**
    - defining threshold for respiratory depression, 41-42
    - intravenous patient-controlled analgesia with opioid, 40-41
    - practical considerations for, 44-45
    - predicting settings for, 42-43
    - transition to oral opioids, 46
    - opioid tolerance, 38
    - preoperative fentanyl challenge, 41
    - preoperative plan for, 38-39
    - regional analgesia techniques and nonopioid analgesics, 39-40
    - epidural analgesia and peripheral nerve blocks, 39
    - nonopioid analgesics, 39-40
  - inadequate, consequences of, **21-36**
    - for recovery and health care use, 24-25
    - persistent postoperative pain, 25-31
      - definition, 25
      - incidence, 25-26
      - mechanisms involved in, 28-29
      - predictors of, 26-28
      - prevention of, 29-31
    - physiologic, 21-23
    - psychologic and behavioral, 23-24
  - incisional pain, mechanisms of, **1-20**
    - clinical models of, 2-4
    - experimental *versus* clinical models of, 4

- laparotomy models for, 4-5
  - plantar incision model for, 5-15
  - intravenous patient-controlled analgesia, **109-123**
  - local anesthetics, clinical pharmacology of, **73-84**
    - beneficial actions, 79-81
    - long-lasting, 77-78
    - novel delivery techniques, 75-77
      - catheter techniques, 76-77
      - encapsulated drugs, 75-76
    - shift to outpatient surgery and, 74-75
    - toxicity issues, 78-79
    - unexpected toxicity, 73-74
  - multimodal analgesia techniques and rehabilitation, **185-202**
    - analgesic options for, 186-193
      - as part of preemptive analgesia techniques, 195-196
      - optimal techniques for, 193-195
      - to shorten convalescence, 196
  - peripheral nerve blocks and continuous catheter techniques, **141-162**
    - brachial plexus blocks, 141-146
      - axillary, 145-146
      - infraclavicular, 144-145
      - interscalene, 141-143
      - supraclavicular, 143-144
    - femoral plexus block, 148-150
    - lumbar plexus block, 146-148
    - paravertebral nerve block, 153-154
    - popliteal fossa sciatic nerve block, 151-153
    - sciatic nerve block, 150-151
  - procedure-specific, **203-210**
    - example of, for laparoscopic cholecystectomy, 207-209
    - initiatives to provide guidelines for, 205-209
    - reasons for, 204-205
- Pain scales, for children, 164
- Pain scores, defining maximum acceptable in acute pain service, 219
- Pain, postoperative, incisional, mechanisms of, **1-20**
  - clinical models of, 2-4
  - experimental *versus* clinical models of, 4
  - laparotomy models for, 4-5
  - plantar incision model for, 5-15
  - prevention strategies, 15-16
  - schematic for, 16-17
  - persistent, as consequence of inadequate pain relief, 25-31
    - definition, 25
    - incidence, 25-26
    - mechanisms involved in, 28-29
    - predictors of, 26-28
    - prevention of, 29-31
- Palliative care, in pediatrics, **837-856**
  - barriers to access to, 851-853
  - epidemiology of life-threatening illnesses, 837-839
  - ethics and decision making, 849-851
    - advance care planning, 849-850
    - Baby Doe, 850
    - in adolescents, 851
  - pain and symptom management, 844-849
  - psychological and spiritual concerns, 839-844
- Paravertebral nerve blocks, for postoperative pain management, 153-154
  - in multimodal approach to, 187-188
- Parecoxib, efficacy in postoperative pain management, 56-57
- Parents, of pediatric patients, preoperative preparation programs for, 600
  - presence during induction of anesthesia, impact on child's anxiety, 601-604
- Parvovirus infection, transfusion-transmitted, 245
- Pathophysiology, in obesity, **421-429**
  - cancer, 427
  - cardiovascular, 422
  - metabolic syndrome, 423-425
  - respiratory, 425-427
  - of obstructive sleep apnea, **431-443**, 494-495
    - categories of, 431-433
    - causes, 434-435
    - contributing factors, 435-436
    - diseases associated with, 438-440
    - in children, 535-537
    - symptoms, 436-437



- Patient-controlled analgesia, intravenous, **109-123**  
 adjuvant drugs and, **112-114**  
   added to the IV-PCA solution, **113-114**  
   administered separately, **112-113**  
 choices of opioids for, **111-112**  
 efficacy of, **109-111**  
 epidural, in children, **175-176**  
 in children, **168-170**  
   epidural, **175-176**  
 in specific patient populations, **117-119**  
   elderly patients, **118-119**  
   morbidly obese patients, **117-118**  
   opioid-tolerant patients, **117**  
   sleep apnea patients, **117-118**  
 opioid-related side effects, **112**  
 patient factors, **116-117**  
   patient education, **116-117**  
   psychological characteristics, **116**  
 pediatric acute pain management with, **801-803**  
   epidural, **805-806**  
 program parameters for, **114-116**  
   bolus dose, **114-115**  
   concurrent background (continuous) infusions, **115-116**  
   dose limits, **116**  
   lockout interval, **115**  
 with opioids, in chronic opioid-consuming patients, **40-46**  
   defining threshold for respiratory depression, **41-42**  
   practical considerations for, **44-45**  
   predicting settings for, **42-44**  
   preoperative fentanyl challenge, **41**  
   transition to oral opioids, **46**
- Pediatric anesthesia, **573-861**  
 acute pain management, **789-814**  
 blood conservation in, **347-361**  
   acute normovolemic hemodilution, **350-353**  
   deliberate hypotension, **354**  
   human recombinant erythropoietin, preoperative administration of, **354-356**  
   intraoperative blood salvage, **353**  
   multimodal strategies for, **356-357**  
   preoperative autologous donation, **349-350**  
   transfusion triggers and institutional algorithms, **348-**  
   cardiac, new developments in, **655-676**  
   for children with cerebral palsy, **733-743**  
   for obese children, **745-764**  
   for spinal fusion, **765-787**  
   for thoracic solid organ transplant, **709-731**  
   future of, **857-861**  
   genetics of malignant hyperthermia, **615-619**  
   historical perspective, **573-595**  
   opioids in, **621-634**  
   palliative care and, **837-856**  
   preoperative psychological preparation for surgery, **597-614**  
   procedural sedation in, **635-654**  
   regional, **815-835**  
   single-lung ventilation in, **693-708**  
   ventilators for, **677-691**
- Pediatrics, obstructive sleep apnea in children, **535-549**  
 clinical features, **537-538**  
 complications, **538**  
 evaluation, **539**  
 pathophysiology, **535-537**  
   airway anatomy, **536**  
   neuromotor factors, **537**  
   obesity and, **536-537**  
 treatment, **539-547**  
   intraoperative anesthesia plan, **542-544**  
   monitoring, **544-545**  
   pain management, **546**  
   perioperative management, **540**  
   postoperative care, **544**  
   preanesthesia evaluation, **540-542**  
   prognosis, **546-547**  
   respiratory support, **545-546**  
   surgery for, **540**  
 postoperative pain management in children, **163-184**  
   epidural/caudal blocks, **170-175**  
   intravenous patient/parent/nurse-controlled analgesia, **168-170**  
   local anesthetics/adjuncts, **176-180**  
   nonopioid analgesics, **165-168**  
   potent opioid analgesics, **168**
- Penile nerve block, in children, **818-819**
- Perflubron emulsion (Oxygent), **382-384**
- Perfluorocarbon (PFC) emulsions, as artificial oxygen carriers, **382-384**
- Perioperative assessment, of elective orthopedic surgery patient, **295-303**
- Perioperative care, for surgical treatment of obstructive sleep apnea, **531-532**  
 in children, **540**
- Peripheral nerve blocks, for pediatric acute pain management, **808-809**  
 for postoperative pain, **141-162**  
   brachial plexus blocks, **141-146**  
   axillary, **145-146**  
   infraclavicular, **144-145**

- interscalene, 141–143
  - supraclavicular, 143–144
  - femoral plexus block, 148–150
  - in chronic opioid-consuming patients, 39
  - lumbar plexus block, 146–148
  - paravertebral nerve block, 153–154
  - popliteal fossa sciatic nerve block, 151–153
  - sciatic nerve block, 150–151
  - in multimodal approach to postoperative pain management, 187–188
- Persistent postoperative pain. *See Pain, postoperative, persistent.*
- Pharmacokinetics, anesthetic management in obese patients, 479–481
- Pharmacology, developmental, and opioids in pediatric anesthesia, 622–625
- in obese children and adolescents, 754–756
  - of agents for procedural sedation in pediatric patients, 644–650
  - of induction agents, and rapid sequence induction in patients with obesity and/or obstructive sleep apnea, 560–561
  - of pain behaviors after plantar incision, 8–9
  - spinal and parenteral, of incisional pain, 12–15
- Pharmacotherapy, in management of obesity, 502–504
- Phen-fen, 503
- Phentermine, 503
- Phlebotomy, aggressive, in autologous blood donation, 266–267
- Physiologic consequences, of inadequate postoperative pain relief, 21–23
- Physiotherapy, of lingual and supra-hyoid muscles, for treatment of obstructive sleep apnea, 531
- Plantar incision model, for postoperative pain, 5–15
- dorsal horn neuron sensitization, 12
  - nonevoked pain, 5–7
  - primary afferent fiber sensitization, 10–12
  - primary mechanical hyperalgesia, 7, 10
  - spinal and parenteral pharmacology of, 12–15
- Polymerized hemoglobin-based oxygen carrier, bovine, 378–379
- human, 379–380
- Polysomnography, in diagnosis of obstructive sleep apnea, 406–408
- in children, 539
- Popliteal fossa sciatic nerve block, for postoperative pain, 151–153
- Positioning, for surgery, obese patients, 481–482
- Post-transplant lymphoproliferative disorder, after pediatric lung transplantation, 726–727
- Postanesthesia care unit (PACU), management of patients with obstructive sleep apnea in, 495–497
- disposition from, 496–497
  - during initial recovery, 495–496
- Postoperative management, of patients with obstructive sleep apnea, **493–500**
- additional considerations for obese patients, 498
  - children, 544
  - disposition from postanesthesia care unit, 496–497
  - during recovery and immediate postoperative period, 495–496
  - pain management, 497–498
- Postoperative pain, models of, clinical, 2–4
- experimental *versus* clinical, 4
  - laparotomy, 4–5
  - plantar incision, 5–15
- persistent, as consequence of inadequate pain relief, 25–31
- definition, 25
  - incidence, 25–26
  - mechanisms involved in, 28–29
  - predictors of, 26–28
  - prevention of, 29–31
- Postoperative pain management. *See Pain management.*
- Preemptive analgesia, multimodal analgesia as part of, 195–196
- Pregabalin, in multimodal approach to postoperative pain management, 193
- Preoperative autologous blood donation, **263–270**
- aggressive phlebotomy in, 266–267
  - cost effectiveness, 268
  - erythropoietin use in, 266–267
  - for elective orthopedic surgery, perioperative assessment of patient, **295–303**
  - preoperative evaluation and methods to reduce blood use, **305–313**

- in pediatric patients, 349–350
  - patient selection for, 264–266
  - preoperative collection of components, 267–268
  - transfusion trigger in, 267
  - Preoperative evaluation, and methods to reduce blood use in orthopedic surgery, **305–313**
    - of patients with obesity, **463–469**
      - airway assessment, 465–466
      - cardiovascular system assessment, 468–469
      - other considerations, 469–470
      - respiratory function assessment, 466–467
    - of patients with obstructive sleep apnea, **470–474**
      - assessment of patient with, 472–473
      - cardiovascular assessment, 473–474
      - identifying patients with, 470–472
      - in children, 540–542
  - Preoperative preparation, psychological, of child for surgery, **597–614**
    - risk factors for preoperative anxiety, 598–599
    - treatment of preoperative anxiety, 599–606
      - acupuncture, 605–606
      - behavioral intervention, 606
      - music, 604–605
      - outcomes of, 606–609
      - parental presence during induction of anesthesia, 601–604
      - preoperative preparation programs, 599–601
  - Prescription errors, in transfusion medicine, 256
  - Pressure controlled ventilation, in pediatric anesthesia, 685–687
  - Pressure support ventilation, in pediatric anesthesia, 688
  - Prevalence, of obstructive sleep apnea, 408–410
  - Procedural sedation, in pediatric patients, **635–654**
  - Procedure-specific postoperative pain management, **203–210**
    - example of, for laparoscopic cholecystectomy, 207–209
    - initiatives to provide guidelines for, 205–209
    - reasons for, 204–205
  - Prognosis, of children with obstructive sleep apnea, 546–547
  - Propofol, for procedural sedation in pediatric patients, 648–649
  - Protein binding, developmental pharmacology and opioids in pediatric anesthesia, 622–625
  - Psychologic consequences, of inadequate postoperative pain relief, 23–24
  - Psychological preparation, preoperative, of child for surgery, **597–614**
    - risk factors for preoperative anxiety, 598–599
    - treatment of preoperative anxiety, 599–606
      - acupuncture, 605–606
      - behavioral intervention, 606
      - music, 604–605
      - outcomes of, 606–609
      - parental presence during induction of anesthesia, 601–604
      - preoperative preparation programs, 599–601
  - Pulmonary aspiration, risk of, in anesthesia of patients with obesity and/or obstructive sleep apnea, 555–556
  - Pulmonary circulation, modulation of, in single-lung ventilation for pediatric anesthesia, 705
  - Pulmonary hypertension, in patients with obstructive sleep apnea, preoperative evaluation of, 473
- 
- R**
- Race, as risk factor for obstructive sleep apnea, 413
  - Racial differences, in obesity, 398, 401
  - Radiofrequency tissue ablation, multilevel, for treatment of obstructive sleep apnea, 531
  - Rapid sequence induction, in patients with obesity and/or obstructive sleep apnea, **551–564**
    - cricoid pressure, 556–560
    - induction agent pharmacology, 560–561
    - risk of gastroesophageal reflux and pulmonary aspiration, 555–556
    - with the difficult airway, 552–555
  - Recombinant factor VIIa, in pediatric cardiac anesthesia, 660–661

- Recombinant human erythropoietin, blood conservation with, in cardiac surgery, 339-340  
in elective orthopedic surgery, 309-311  
in pediatric surgical patients, 354-356
- Recovery, of red blood cells. *See Cell salvage.*
- Red blood cell recovery and reinfusion. *See Cell salvage.*
- Regional analgesia, for postoperative pain in chronic opioid-consuming patients, 39
- Regional anesthesia, in children, **815-835**  
axillary block/brachial plexus blocks, 822-824  
caudal and lumbar, 824-827  
epidural opiates, 828-830  
fascia iliaca compartment block, 821  
femoral nerve block, 819-821  
ilioinguinal-iliohypogastric nerve block, 816-818  
local infiltration and "splash" local anesthesia, 824  
penile nerve block, 818-819  
sciatic nerve blockade, 821-822  
spinal, for infants, 830-832  
volume and dosage of local, 827-828  
in obese patients, 482-483
- Regional techniques, for pediatric acute pain management, 807
- Rehabilitation, postoperative, multimodal analgesia to shorten, 196
- Reinfusion, of recovered red blood cells. *See Cell salvage.*
- Rejection, after pediatric solid organ transplantation, heart, 715  
lung, 724
- Remifentanyl, in pediatric anesthesia, 628-629  
new developments in pediatric cardiac anesthesia, 656-657
- Renal drug clearance, developmental pharmacology and opioids in pediatric anesthesia, 622-625
- Reservoir, collection, in red cell recovery and reinfusion, 287-288
- Respiratory depression, predicting threshold for, with postoperative opioids in chronic opioid-consuming patients, 41-42
- Respiratory function, preoperative assessment in obese patients, 466-467
- Respiratory pathophysiology, in obesity, 425-427
- Respiratory support, after surgery in children with obstructive sleep apnea, 545-546
- Risk factors, for obesity, **397-403**  
for obstructive sleep apnea, 410-415  
age, 412-413  
alcohol consumption, 414-415  
craniofacial abnormalities, 414  
family history, 414  
gender, 412  
obesity, 411-412  
race, 413
- Ropivacaine, for postoperative pain in children, 177-178
- Roux-en-Y gastric bypass, surgical management of obesity with, 511-514  
complications following, 513  
essential components of, 512
- 
- S**
- Salvage. *See Cell salvage.*
- Sampling, pretransfusion, errors in, 256-257
- Sciatic nerve block, for postoperative pain, 150-151
- Sciatic nerve blockade, in children, 821-822
- Sedation, procedural, in pediatric patients, **635-654**  
controversial issues in, 650-651  
documentation, 642  
equipment for, 639-641  
goals of, 636  
levels of, 636-637  
monitoring, 641-642  
patient selection, 638-639  
pharmacology of, 644-650  
presedation planning and consent, 639  
recovery and discharge, 642-644  
skills and training of personnel for, 637-638
- Sensitization, dorsal horn neuron, 12  
hyperalgesia and, 2  
primary afferent fiber, 10-12
- Serious Hazards of Transfusion (SHOT) scheme, 254-255
- Silbutramine, in management of obesity, 503-504
- Single-lung ventilation, in pediatric anesthesia for thoracic surgery, **693-708**  
avoiding airway injury during, 704  
modulation of pulmonary circulation, 705-706  
monitoring and anesthetic techniques, 704-705

- pain management techniques, 705  
 physiology of, 695-696  
     lateral decubitus position, 693-695  
 surgical lesions of chest, 696  
 techniques, 697-703  
     Arndt endobronchial blockers, 698-700  
     balloon-tipped bronchial blockers, 697  
     double-lumen tubes, 701-703  
     Fogarty embolectomy catheters, 697-698  
     in patients who have a tracheostomy, 703  
     single-lumen endotracheal tube, 697  
     Univent tubes, 700-701  
     treating and avoiding hypoxemia during, 703-704  
     video-assisted thoracoscopic surgery, 705
- Sleep apnea. *See Obstructive sleep apnea.*
- Sleep regulation, abnormalities in, inherited, predisposing to sleep apnea, 456-457
- Sleep study. *See Polysomnography.*
- Socioeconomic factors, role in obesity, 401
- Sodium bicarbonate, as adjunct for postoperative pain in children, 179
- Solid organ transplant surgery, pediatric, anesthesia considerations for, **709-731**  
     heart transplantation, 709-715  
     lung transplantation, 715-728
- Spinal anesthesia, for infants, 830-832  
     anatomy, 830-831  
     drug dosage in, 832  
     technique in ex-premature patients, 831
- Spinal fusion, pediatric, anesthetic management for, **765-787**  
     anesthesia, 774-781  
         clinical synthesis, 780-781  
         inhaled anesthetics, 775-776  
         intravenous anesthetics, 776-779  
         nonanesthetic intraoperative influences, 779  
     evoked potentials, 767-774  
     need for intraoperative neurologic monitoring, 766-767  
     preoperative evaluation, 765-766
- Spiritual care, as part of pediatric palliative care, 842
- Splash local anesthesia, in children, 824
- Sponges, surgical, rinsing of, in red cell recovery and reinfusion, 286-287
- Staff support, in pediatric palliative care, 843-844
- Stress response, new developments in pediatric cardiac anesthesia, 656
- Stroke, associated with obstructive sleep apnea, 439
- Sufentanil, in pediatric anesthesia, 627-628
- Supra-hyoid muscles, physiotherapy of, for treatment of obstructive sleep apnea, 531
- Supraclavicular brachial plexus block, for postoperative pain, 143-144
- Surgeons, role in acute pain service, 219
- Surgery, for children who have cerebral palsy, **733-743**  
     for obstructive sleep apnea, 527-532  
         jaw advancement techniques, 530  
         nasal surgery, 529  
         perioperative issues for, 531-532  
         physiotherapy of lingual and supra-hyoid muscles, 531  
         tonsillectomy, 528-529  
         tracheostomy, 529-530  
         uvulopharyngopalatoplasty, 527-528  
     for weight loss. *See Bariatric surgery.*  
     positioning, with obese patients, 481-482  
     preoperative psychological preparation of child for, **597-614**  
     thoracic, single-lung ventilation for children undergoing, **693-708**
- Synchronized intermittent mandatory ventilation, in pediatric anesthesia, 687

---

**T**

- Thoracic epidural analgesia, continuous, in children, 174-175
- Thoracic surgery, pediatric solid organ transplant, anesthesia considerations for, **709-731**  
     heart transplantation, 709-715  
     lung transplantation, 715-728  
     physiology of lateral decubitus position, 693-695  
     single-lung ventilation in pediatric anesthesia for, **693-708**  
     surgical lesions of chest, 696  
     video-assisted, 705

- Tissue ablation, multilevel radiofrequency, for treatment of obstructive sleep apnea, 531
- Tolerance, opioid, 38, 117
- Tongue-base suspension, for treatment of obstructive sleep apnea, 531
- Tonsillectomy, for treatment of obstructive sleep apnea, 528-529  
in children, for obstructive sleep apnea treatment, 539-540
- Toxicity, of local anesthetics in postoperative pain management, 78-79
- Tracheostomy, for treatment of obstructive sleep apnea, 529-530  
single-lung ventilation for pediatric anesthesia in child with, 703
- Tramadol, as adjunct for postoperative pain in children, 180  
for postoperative pain in children, 167-168  
pediatric acute pain management with, 797
- Transcranial Doppler, in pediatric cardiac anesthesia, 663-664
- Transfusion medicine, 241-389  
acute normovolemic hemodilution, 271-281  
benefits of, 277  
cost effectiveness of, 278  
efficacy of, 277-278  
future of, 278  
in pediatric patients, 350-353  
physiologic consequences, 273-274  
practical considerations for, 275-277  
technique, 271-273  
anemia and, in surgical patients, 315-325  
efficacy of transfusion, 319-322  
human data, 318-319  
interaction between pathophysiologic processes and, 317-318  
pathophysiology, 315-317  
transfusion threshold, 322-323  
artificial oxygen carriers, 373-389  
autologous blood donation, 263-270  
aggressive phlebotomy in, 266-267  
cost effectiveness, 268  
erythropoietin use in, 266-267  
patient selection for, 264-266  
preoperative collection of components, 267-268  
transfusion trigger in, 267  
blood conservation, in critically ill patients, 363-372  
in elective orthopedic surgery, 305-313  
in pediatric anesthesia, 347-361  
cardiac surgery, conventional *versus* bloodless, 327-345  
blood conservation methods in, 339-341  
clinical outcomes, factors influencing, 334-339  
coronary artery bypass graft techniques and, 330  
laboratory coagulation tests, 330-331  
predictors of transfusion requirements for, 332-334  
variability in transfusion practice for, 331-332  
errors in, 247-248, 253-261  
prevention of, 258-259  
types of, 255-258  
UK Serious Hazards of Transfusion (SHOT) scheme, 254-255  
in critically ill patients, 363-372  
in elective orthopedic surgery, perioperative patient assessment, 295-303  
preoperative evaluation and reducing blood use in, 305-313  
in pediatric patients, 347-361  
red blood cell recovery and reinfusion, 283-294  
comparison with other techniques, 290-292  
cell salvage, 292  
normovolemic hemodilution, 291-292  
preoperative autologous donation, 291  
complications, 290  
air embolism, 290  
wrong wash solution, 290  
optimizing red cell return, anticoagulant, 287  
calculation of blood loss during cell salvage, 288  
collection reservoir, 287-288  
optimizing red cell return, 285-288  
rinsing of surgical sponges, 286  
suction, 285-286  
wash quality, 289-290  
physics of cell salvage machine, 283-285  
risks of blood transfusion, 241-252  
acute lung injury related to, 247  
emerging risks, 248  
errors in, 247-248, 253-261  
immunosuppression, 246-247  
infections, 241-246

- Transfusion threshold, increased, to conserve blood, 340-341
- Transfusion triggers, in autologous blood donation, 267  
in pediatric anesthesia, 348
- Transfusion-related acute lung injury, 247
- Transplant surgery, pediatric solid organ transplant, anesthesia considerations for, **709-731**  
heart transplantation, 709-715  
lung transplantation, 715-728
- Trigger, transfusion, in autologous blood donation, 267  
in pediatric anesthesia, 348

---

**U**

- Univent tubes, for single-lung ventilation in pediatric anesthesia, 700-701
- Upper airway resistance. See also *Obstructive sleep apnea*.  
physiologic characteristics of, 431-433
- Uvulopharyngopalatoplasty, for treatment of obstructive sleep apnea, 527-528

---

**V**

- Valdecobix, efficacy in postoperative pain management, 55-56
- Vascular complications, after pediatric lung transplantation, 725
- Ventilation control, alterations in, inherited, predisposing to sleep apnea, 455-456
- Ventilatory support, mechanical, anesthesia ventilators, better options for children, **677-691**  
comparison studies, 688-690  
conventional, 679-681  
modern, 681-685  
selecting the ventilation mode, 685-688  
pressure controlled ventilation, 685-687

- pressure support ventilation, 688
- synchronized intermittent mandatory ventilation, 687
- volume control ventilation, 685
- in pediatric cardiac anesthesia, advances in, 666-667
- single-lung ventilation in pediatric anesthesia for thoracic surgery, **693-708**

- Vertical banded gastroplasty, surgical management of obesity with, 508-509
- Video-assisted thoracoscopic surgery, single-lung ventilation for pediatric anesthesia with, 705
- Viral infections, due to blood transfusions, 242-243  
West Nile virus, emerging risk of, 248
- Volume control ventilation, in pediatric anesthesia, 685

---

**W**

- Weight loss, for treatment of obstructive sleep apnea, 527
- Weight loss drugs, in management of obesity, 502-504
- Weight loss surgery. See *Bariatric surgery*.
- West Nile virus, transfusion-associated, emerging risk of, 248
- Wound healing, effects of cyclooxygenase-2 inhibitors for postoperative pain management, 61-62

---

**X**

- Xenical. See *Orlistat*.